

Diversity, **Minorities and Granovetter's** Model

Universität Konstanz



UNIVERSITÄT KONSTANZ

Computational Modelling of Social Systems

Giordano De Marzo

Outline

1. Chaos and Randomness 2. Collective Behavior and Complex Systems 3. Granovetter's Threshold Model 4. Analysis of Granovetter's Threshold Model 5. Bonus: Can a Minority Win?





Deterministic Chaos



Chaotic is not random! We model the evolution of a population of animals or bacteria with the Logistic

Map:

- $\dot{x}(t)$ = ratio of existing population to the maximum possible population $x(t+1) = r \cdot x(t) \cdot [1-x(t)]$
- 0<r<4 is the parameter of the model Two effects:
 - size is small the population grows is large the population decreases
 - reproduction when the population • starvation when the population size

This model is completely deterministic!

Example of Evolution















Example of Evolution









- The behavior is regular
- Two moderately far initial points results in very similar trajectories

Chaotic Trajectories



- The behavior is apparently randomic

• Two very close initial points results in completely different trajectories https://www.complexity-explorables.org/flongs/logistic/

- the x axis gives x(t)
- the y axis gives x(t+1)
- the straight line is x(t+1)=x(t)





- the x axis gives x(t)
- the y axis gives x(t+1)
- the straight line is x(t+1)=x(t)

The recipe for evolving the system is very simple 1.start from *x(0)* and move up until you intersect the curve



- the x axis gives x(t)
- the y axis gives x(t+1)
- the straight line is x(t+1)=x(t)

The recipe for evolving the system is very simple

1. start from *x(0)* and move up until you intersect the curve

2.move horizontally until you intersect the straight line in x(1)



- the x axis gives x(t)
- the y axis gives x(t+1)
- the straight line is x(t+1)=x(t)

The recipe for evolving the system is very simple

- 1. start from *x(0)* and move up until you intersect the curve
- 2.move horizontally until you intersect the straight line in x(1)3.move vertically until you intersect

the curve again

1.0 -0.9 0.8 0.7 -0.6 x(t+1)0.5 -0.4 0.3 0.2 -0.1



- the x axis gives x(t)
- the y axis gives x(t+1)
- the straight line is x(t+1)=x(t)

The recipe for evolving the system is very simple

- 1. start from x(0) and move up until you intersect the curve
- 2. move horizontally until you intersect the straight line in x(1)
- 3. move vertically until you intersect the curve again
- 4. repeat from step 2

1.0 -

0.9

0.8 -

0.7 -

0.6 -

0.5 -

0.4

0.3

0.2 -

0.1

x(t+1)



- the x axis gives x(t)
- the y axis gives x(t+1)
- the straight line is x(t+1)=x(t)

The recipe for evolving the system is very simple

- 1. start from x(0) and move up until you intersect the curve
- 2. move horizontally until you intersect the straight line in x(1)
- 3. move vertically until you intersect the curve again
- 4. repeat from step 2

1.0 -

0.9

0.8 -

0.7 -

0.6 -

0.5 -

0.4

0.3

0.2 -

0.1

x(t+1)



- the x axis gives x(t)
- the y axis gives x(t+1)
- the straight line is x(t+1)=x(t)

The recipe for evolving the system is very simple

- 1. start from x(0) and move up until you intersect the curve
- 2. move horizontally until you intersect the straight line in x(1)
- 3. move vertically until you intersect the curve again
- 4. repeat from step 2

1.0 -

0.9

0.8 -

0.7 -

0.6 -

0.5 -

0.4

0.3

0.2 -

0.1

x(t+1)



Periodic and Chaotic Trajectories



https://www.complexity-explorables.org/flongs/logistic/

Chaos (Period = ∞)





More is Different!



"The ability to reduce everything to simple fundamental laws does not imply the ability to start from those laws and reconstruct the universe. [...] Psychology is not applied biology, nor is biology applied chemistry. [...] At each level of complexity entirely new properties appear." – Philip Anderson

More is different: broken symmetry and the nature of the hierarchical structure of science. Philip Anderson, Science (1972)

Complicated or Complex?





Complicated System

Example: a mechanical watch

- pieces have specific functions and welldefined relationships
- carefully engineered or designed
- it is easy to infer global behavior and understand outcomes of modifications

Complex System

Example: a human cell

- pieces have unknown functions and relationships
- understand outcomes of modifications
- Self-organized, no external project • it is hard to infer global behavior and

Jurassic Park, Chaos and Complexity

Jurassic Park is not a book about dinosaurs, it is a book about chaos and complex systems!

"Chaos theory throws it right out the window. It says that you can never predict certain phenomena at all. You can never predict the weather more than a few days away. All the money that has been spent on long-range forecasting-about half a billion dollars in the last few decades-is money wasted. It's a fool's errand. It's as pointless as trying to turn lead into gold." – Dr Ian Malcolm, Jurassic Park



Collective Behavior Once Again

Complex Systems are characterized by emergent collective behaviors.

Nature is full of collective behavior examples:

- flocks of bird
- schools of fish
- ants and bees

Collective behavior can emerge even in very simple animals!



Diversity-Induced Collective Behavior

Interaction-induced collective behavior

The macro behavior depends on the **interaction** between individuals:

- Schelling's model: low tolerance triggers moves that lead to segregation
- Alxelrod's model: cultural exchange leads to larger cultures or supports coexistence of few cultures

Diversity-induced collective behavior

The macro behavior emerges from **differences** between individuals. Same interaction pattern can lead to very different outcomes

veen individuals: at lead to segregation cultures or supports



The Riot Toy Example

A group of individuals is part of a demonstration:

- Individuals have a **threshold** of how many others have to be rioting to join the riot
- If enough people are in the riot, individuals with lower threshold join too

This is an example of **binary opinions**

- Proto-opinion: just participate / not participate
- Other examples with binary decisions depending on size: Diffusion of innovations, rumors, strikes, voting...



An Example of Spreading



Video unavailable Watch on YouTube



Rational Agents in Collective Actions

Cost

Benefit

We assume agents to be **rational**, so the decision to join the collective action depends on: • **Risk** or **cost** of participating. • Risk of being jailed in riot • Wage loss in strike Cost of technology adoption • The **benefit** (potential or sure) of the action taking place. • Political change after demonstration Political party winning an election Profit out of adopting

innovation



Net benefit = benefit - costs
Threshold to join: Net benefit must be positive (>0)
benefits increase and costs

- decrease with more people in the action
 - group effect on social network
 - less possibility to be arrosted in riot
 - arrested in riot
 - economy of scale following
 tochnology adoption
- technology adoptionweaker assumption: there is
 - only one crossing of zero in the function of net benefit vs people
 - in action

Questions on Spreading and Diversity

We want to understand the role of **Diversity** in inducing the spreading of ideas or behaviors

- How does the distribution of preferences (thresholds) in a population affect its collective behavior?
- Knowing the preferences does not directly tell you how the population will behave, you need to analyze how the population behaves
- Aim: understanding groups beyond the representative "mean" member



- Granovetter's model schematizes the process of joining a riot • there are *n* distinct agents
 - each agent is characterized by a threshold θ_i
 - an agent join the riot if and only if there is a number of agents larger or equal to its threshold in the riot • we denote by **M(t)** the number of rioters at time t (and by **x(t)** the
 - percentage)
 - at the time step *t*+1 all agents with $\theta_i \leq M(t)$ join the riot
 - the simulation stops if all people are in the riot M(t) = n or a stationary state M(t+1)=M(t) is reached

One Example with Spreading



- *n* Agents
- Uniform sequence of thresholds with integer values [0, n-1]
- First agent activates, then second, and so on
- One agent joins per iteration and all agents are active in the end

One Example without Spreading



- Same example as before but agent with threshold 1 now has threshold 2
- First agent activates and simulation ends
- Radically different outcome for minimal change in thresholds!
- Deducing preference distributions from collective outcomes is risky

In a real group of people there will be an average behavior with some fluctuations (very violent or very pacific people)

- Thresholds follow normal distribution with mean *µ* and standard deviation σ
- we denote by \mathbf{x}_{e} the equilibrium percentage of active agents
- Number of agent is *N=100*
- Mean value is constant $\mu = 25$
- Sharp increase in x_e at a critical σ value: discontinuous or first order phase transition
- Diversity-induced collective behavior







standard deviation of thresholds o



Granovetter's Model as a Map

Granovetter's threshold model is just a map!

- x(t) is the fraction of rioters at time t
- x(t+1) only depend on the thresholds (fixed) and x(t)
- the process is deterministic
- we have to understand the function that makes x(t)evolve into x(t+1)

rioters is

where $N[\theta \le M(t)]$ is the number of agents with threshold less than M(t). We divide both size by the number of agents n

 $x(t+1)=N[\theta \leq M(t)]/n \approx P[\theta \leq M(t)]$ $P[\theta \le M(t)]$ is the cumulative probability of the thresholds. For simplifying things we normalize the thresholds $\Theta = \theta/n$ and we obtain

The evolution of the number of

 $M(t+1)=N[\theta \leq M(t)]$

 $x(t+1) \approx P[\Theta \leq x(t)]$

Cumulative Probability 0.5

0.5

- P(x) =
- - P(x) =

The figure shows the cumulative of a gaussian

The cumulative probability P(x)is the probability to extract a number smaller than a given value x. If p(x) is the probability to extract a number x then for discrete variables

=
$$\sum_{y=0}^{x} p(y)$$

for continuous variables

$$=\int_0^x p(y)dy$$



• the map has 3 equilibrium points

- *x~0* stable
- *x~0.14* unstable
- \circ *x=1* stable
- since x(0)=0 the stable point
 - x=1 is never reached
- around 0% of the agents
 - involved in the riot



• the map has 3 equilibrium points

- *x~0.04* stable
- *x~0.14* unstable
- \circ *x=1* stable
- since x(0)=0 the stable point
 - x=1 is never reached
- around 4% of the agents
 - involved in the riot



• the map now has only 1 equilibrium point \circ *x=1* stable • starting from x(0)=0 the system reaches the only stable equilibrium point • 100% of the agents involved in the riot



- the map now has only 1 equilibrium point • *x=0.83* stable • starting from x(0)=0 the system reaches the only stable equilibrium point • 83% of the agents involved in the riot

- there is a decrease for large
 - variance

Take Home Messages

Modelling action as rational choice

Thresholds as points where benefits outweigh costs or risks **Diversity matters**

Two populations with the same average threshold have very different behaviors even if mean thresholds are the same

Tipping point or phase transition

Behavior changes dramatically at a narrow range of standard deviation of thresholds

Size effects

Small changes in threshold sequences can be important. When the population is small, you have a probability of very different outcomes. Inferring the preferences from the outcome is very hard and/or misleading



What is a Minority?



- Often based on
 - ethnicity
 - religion
 - sexual orientation
 - gender
- Not necessarily numerical



Minority groups are categories of people differentiated from the social majority

• They face inequalities and discrimination • Play a critical role in social movements and in initiating changes in societal norms

Salary Gap

The unadjusted gender pay gap, 2022

(difference between average gross hourly earnings of male and female employees as % of male gross earnings)



Note: For all the countries except Czechia and Iceland: data for enterprises employing 10 or more employees, NACE Rev. 2 B to S (-O); Czechia: data for enterprises employing 1 or more employees, NACE Rev. 2 B to S; Iceland: NACE Rev. 2 sections C to H, J, K, P, Q. Gender pay gap data for 2022 are provisional until benchmark figures, taken from the Structure of Earnings survey, become available in December 2024.

* Euro area (2015-2022)

(1) Estimated data.

(2) Definition differs (see metadata). (3) 2018 data.

Source: Eurostat (online data code: sdg_05_20)





Never Stop at the First Graph!

Employment rate of women and men in the EU (as % of the population aged 20 to 64, 2018 data)



ec.europa.eu/eurostat



Stubborn Minorities



A stubborn minority is a minority that will never change

its habits, no matter what:

- Jewish and Kosher food
- Sexual habits
- Religion

When the majority has no interest in the specific matter, a change in the social norms may occur. The idea is that it is easier to have all Kosher beverages than having to produce and distribute two different products.

The Most Intolerant Wins: The Dictatorship of the Small Minority Nassim Nicholas Taleb

Critical Mass Theory

Apparently stable societal norms can be effectively overturned by the efforts of small but committed minorities. This leads to the Critical Mass Theory

- when a committed minority reaches a critical group size the social system crosses a tipping point
- Once the tipping point is reached, the actions of a minority group trigger a cascade of behavior change



Modeling the Tipping Point

We consider a model similar to the Naming Game

Agent

Agents store the last M names (or strategies) they heard



Space

Agents interact on a fully connected network (mean field)



Dynamics

The speaker communicates the most common word in its memory. The speaker records it.



Committed Agents

We want to study if a committed minority can change social norms

- we set the initial state with all memories full of the name A (consensus)
- we introduce stubborn or committed agents
- committed agents always communicate name B, independently how their memory

d age







Simulation Results

The model show a tipping point for a fraction of committed agent around 20/30%. The exact tipping point depends on the memory length.



Testing the Model

MIT/Penn | Name Game OWelcome, nargle



194 Participants divided into 10 independent online groups **Procedure:** Participants randomly paired in rounds within their groups to name a pictured object The objective was to coordinate on the same name with their partner Successful coordination -> financial

- reward

Goal and Incentives:

The aim was not to achieve a global consensus but to coordinate successfully in each pairwise interaction. **Over time a common name emerges!**

• Failure -> financial penalty.

Adding a Committed Minority

After establishing a convention among all participants, a small number of confederates, termed as "committed minority," is introduced into each group:

- Their role was to challenge and attempt to change the established naming convention by consistently using a novel alternative (stubborn)
- The size of the committed minority varied across the 10 groups, designed to study the dynamics of how a critical mass can influence social norms.
- Minority sizes ranged from 15% to 35% of each group's population.



When the committed minority overcomes a threshold, there is a shift of the social norm



Tipping Point

The plot shows the aggregated results from the 10 groups:

- tipping point at 25%
- sharp (first order) transition
- a committed minority can overturn a social norm
- results similar to model

| | 100% - | |
|---|--------|-----|
| Adoption at Last Round Non-Committed Players | 75% - | |
| | 50% - | |
| | 25% - | |
| | 0%- | 10% |

Empirical Trials



Take Home Messages

Committed Minority

A minority that will never conform to the social norm independently of the social pressure of the majority

Social Tipping Points

A committed minority can overturn a societal norm producing an abrupt transition in the system. This occur in correspondence to a critical mass. **Modeling Tipping Points**

We can include stubborn or committed agents in a simple agent based model. Results show a norm transition when the minority size is around 25%. **Experimental Results**

The Agent Based Model is replicated using human participant on an online platform. Similar norm transitions are observed when the committed minority size is 25%.

Conclusions

Chaos and Randomness

Chaos exists in deterministic systems, while randomness in nondeterministic ones. **Diversity Induced Collective Behavior**

Complex Systems show collective behavior that originates from the presence of differences among the individuals.

Granovetter's Threshold Model

A model to describe the activation of individuals based on thresholds. Individual differences have very relevant outcomes.

Analysis of Granovetter's Threshold Model

The model can be described as a map that shows a first order transition (tipping point) when the variance of the thresholds distribution is increased. **Can a Minority Win?**

A stubborn minority can generate an abrupt societal norm change when a critical mass is reached. Model and experiment support this theory.

Quiz

- Which of these is complex and which is complicated?
 - An airplane
 - The Internet
 - The Web
 - A deep neural network
- Do you know any example of spreading?
- For a given σ , does μ change the outcome in Granovetter's model?
- Do you think there are minorities in Konstanz? Are they discriminated?
- Which are some examples of stubborn minorities?
- Do you know any example of tipping point in society?
- Which are the limits and strengths of the tipping point model?

Play Yourself to Understand!

Logistic Map

https://www.complexity-explorables.org/flongs/logistic/